Exercises for Lecture Course on Numerical Optimal Control (NOC) Albert-Ludwigs-Universität Freiburg – Summer Term 2020

Exercise 8: Continuous-Time Optimal Control

Prof. Dr. Moritz Diehl, Andrea Zanelli, Dimitris Kouzoupis, Florian Messerer

Consider the following continuous-time optimal control problem:

$$\min_{\substack{x(t), u(t) \\ \text{s.t.}}} \int_{t=0}^{T} L(x(t), u(t)) dt + E(x(T)) \\
\text{s.t.} \quad x(0) = \bar{x}_{0} \\
\dot{x}(t) = f(x(t), u(t)), \quad t \in [0, T].$$
(1)

- 1. (a) Discretize problem (1) using the explicit Euler integrator with step-size h over N intervals. Write on paper the obtained discrete-time optimal control problem.
 - (b) Write the first-order optimality conditions for the discretized problem obtained in (a). Use the Hamiltonian function defined as

$$H(x, u, \lambda) := L(x, u) + \lambda^T f(x, u)$$
⁽²⁾

to simplify these conditions.

- (c) Now let $N \to \infty$ and $h \to 0$. What type of problem do the conditions derived in (b) converge to?
- (d) Fix N = 2 and apply the Newton method to the first-order optimality conditions for the discretized optimal control obtained in (b). Derive the form of the linear systems associated with the Newton steps. Order the variables as $z = (\lambda_0, x_0, u_0, \lambda_1, x_1, u_1, \lambda_2, x_2)$ and the KKT conditions accordingly as $\nabla_z \mathcal{L}(w) = 0$, where $\mathcal{L}(z)$ is the Lagrangian of the NLP.

For notational simplicity we suggest you use the abbreviations $Q_k := h \nabla_x^2 H(x_k, u_k, \lambda_k)$, $R_k := h \nabla_u^2 H(x_k, u_k, \lambda_k)$, $S_k := h \nabla_{ux}^2 H(x_k, u_k, \lambda_k)$, $A_k := I + h \nabla_x f(x_k, u_k)^{\top}$, $B_k := h \nabla_u f(x_k, u_k)^{\top}$ for $k \in \{0, \ldots, N-1\}$ and $Q_N := \nabla_x^2 E(x_N)$

- (e) [Bonus] The linear systems associated with the Newton steps in (d) can be solved exploiting the Riccati Difference Equation (equation 8.5 in the course's script). Derive this equation.
- (f) [Bonus] What kind of matrix ODE does the difference equation derived in (e) converge to for N → ∞ and h → 0?
 Hint: if you have not solved the honus point (e) you can refer to equation 8.5 from the

Hint: if you have not solved the bonus point (e) you can refer to equation 8.5 from the course's script.